MYCOTAXON

Volume 112, pp. 335–338 *April–June 2010*

Additions to the knowledge of aphyllophoroid fungi (*Basidiomycota*) of Atlantic Rain Forest in São Paulo State, Brazil

Adriana de Mello Gugliotta[1*]

Margarida Pereira Fonsêca[2]

Vera Lúcia Ramos Bononi[1]

** agugliottaibot@yahoo.com.br*

[1]Instituto de Botânica, Seção de Micologia e Liquenologia
Caixa Postal 3005, CEP 01061-970, São Paulo, SP, Brazil

[2]Universidade Guarulhos, Ciências Biológicas
Praça Tereza Cristina, 88 CEP 07023-070, Guarulhos, SP, Brazil

Abstract — The list of aphyllophoroid fungi of the Atlantic Rain Forest in the state of São Paulo is updated. Specimens were collected in four different areas of the Atlantic Rain Forest from 1988 to 2007. Exsiccates deposited in the Herbarium SP were also studied. A list of 85 species of *Basidiomycota* distributed into 11 families and four orders (*Agaricales*, *Hymenochaetales*, *Polyporales*, *Russulales*) is presented. All species are mentioned for the first time for the collection sites. Two species are reported for the first time for Brazil and 17 species are recorded for the first time for São Paulo State. The complete list of specimens is available at http://www.mycotaxon.com/resources/weblists.html.

Key words — diversity, macrofungi, neotropics, taxonomy

Introduction

The Atlantic Rain Forest, which has 20,000 species of plants of which 6000 are endemic, is the second largest block of tropical forests of Brazil. This biome, which formerly occupied 1,315,460 km² of Brazilian territory, extending through the region from Osório, Rio Grande do Sul State (29º53'S and 50º16'W) to Cabo de São Roque, Rio Grande do Norte State (05°31'S and 35°16'W), holds today less than 8% of its original extent and has become one the world's top five biological hotspots (Mittermeier et al. 1999, SOS Mata Atlântica/INPE 2009).

The state of São Paulo still holds a significant portion of this important biome (15% of the total remaining forest), largely in protected areas (Secretaria

do Estado de Meio Ambiente 1996, 2000, SOS Mata Atlântica/INPE 2009). It includes several types of tropical ecosystems, such as the coasts of the Atlantic Ocean, the forests of lowlands and slopes of the Serra do Mar, inland forests and woods of *Araucaria* (Secretaria de Estado do Meio Ambiente 1996).

The aim of this study is to contribute to the knowledge of the diversity of aphyllophoroid fungi of the Atlantic Forest and the state of São Paulo, complementing the inventories in the Parque Estadual da Ilha do Cardoso (Bononi 1979a,b,c, 1984, Gugliotta & Capelari 1995, Gugliotta & Bononi 1999) and Parque Estadual das Fontes do Ipiranga (Bononi et al. 1981, Jesus 1993, Soares & Gugliotta 1998, Louza & Gugliotta 2007, Leal & Gugliotta 2008), and expanding the checklist for aphyllophoroid fungi cited from the Brazilian Atlantic Forest by Baltazar & Gibertoni (2009).

Materials and methods

Specimens were collected in four different areas of the Atlantic Rain Forest in the State of São Paulo, from 1988 to 2007:

1. Parque Estadual da Ilha do Cardoso (25°03'S–48°05'W, 22,500 ha), municipality of Cananéia;

2. Reserva Biológica de Paranapiacaba (23°46'S–46°18'W, 336 ha), municipality of Santo André;

3. Parque Estadual das Fontes do Ipiranga (23°39'S–46°37'W, 549.31 ha), municipality of São Paulo;

4. Reserva Florestal da Cidade Universitária "Armando de Salles Oliveira" da Universidade de São Paulo (23°33'S–46°43'W, 10 ha), municipality of São Paulo.

The studied material was deposited in SP herbarium (Holmgren & Holmgren 1998). Exsiccates from these localities deposited in the Herbarium SP were also studied. Micromorphological observations were made from material mounted in 5% KOH and Melzer's reagent; measurements were made in 5% KOH. Nomenclature, taxonomy and author citation followed databases: CBS (http://www.cbs.knaw.nl/databases/) and Index Fungorum (http://www.indexfungorum.org/Names/Names.asp).

Results and discussion

A list of 85 species of *Basidiomycota* distributed into 11 families and four orders (*Agaricales*, *Hymenochaetales*, *Polyporales*, *Russulales*) is presented. *Polyporaceae* is the most represented family with 30 species. All species are mentioned for the first time for the collection sites.

Schizopora trichiliae (Van der Byl) Ryvarden and *Porogramme albocincta* (Cooke & Massee) J. Lowe are recorded for the first time for Brazil.

Seventeen of the identified species are recorded for the first time for São Paulo State: *Amauroderma omphalodes* (Berk.) Torrend, *Coltricia cinnamomea* (Jacq.) Murrill, *Daedalea aethalodes* (Mont.) Rajchenb., *Dichochaete setosa* (Sw.) Parmasto, *Echinoporia aculeifera* (Berk. & M.A. Curtis) Ryvarden, *Henningsia brasiliensis* (Speg.) Speg., *Hymenochaete floridea* Berk. & Broome, *Hymenochaete minuscula* G. Cunn., *Hymenochaete pinnatifida* Burt, *Hymenochaete rubiginosa* (Dicks.) Lév., *Megasporoporia setulosa* (Henn.) Rajchenb., *Nigroporus macroporus* Ryvarden & Iturr., *Perenniporia ohiensis* (Berk.) Ryvarden, *Perenniporia piperis* (Rick) Rajchenb., *Phellinus ferrugineovelutinus* (Henn.) Ryvarden, *Tinctoporellus epimiltinus* (Berk. & Broome) Ryvarden and *Tyromyces fumidiceps* G.F. Atk.

Gugliotta & Capelari (1995) and Gugliotta & Bononi (1999) reported the occurrence of *Trametes versicolor* (L.) Lloyd in Parque Estadual da Ilha do Cardoso, Municipality of Cananéia, but the specimen (SP 193613) was examined later and re-identified as *Coriolopsis caperata* (Berk.) Murrill; the occurrence of *T. versicolor* in this area was not confirmed.

Acknowledgments

We extend our thanks to Clarice Loguercio Leite and Leif Ryvarden who kindly reviewed the manuscript. We are grateful to FAPESP (04/04310-2) for financial support.

Literature cited

Baltazar JM, Gibertoni TB. 2009. A checklist of the aphyllophoroid fungi (*Basidiomycota*) recorded from the Brazilian Atlantic rain forest. Mycotaxon 109: 493–442.

Bononi VL. 1979a. Basidiomicetos do Parque Estadual da Ilha do Cardoso: I. Espécies hidnóides. Rickia 8: 63–74.

Bononi VL. 1979b. Basidiomicetos do Parque Estadual da Ilha do Cardoso: II. *Hymenochaetaceae*. Rickia 8: 85–99.

Bononi VL. 1979c. Basidiomicetos do Parque Estadual da Ilha do Cardoso: III. Espécies clavarióides, teleforóides e estereóides. Rickia 8: 105–121.

Bononi VL. 1984. Basidiomicetos do Parque Estadual da Ilha do Cardoso. IV. Adições às famílias *Hymenochaetaceae*, *Stereaceae* e *Thelephoraceae*. Rickia 11: 43–52.

Bononi VLR, Trufem SFB, Grandi RAP. 1981. Fungos macroscópicos do Parque Estadual das Fontes do Ipiranga, São Paulo, Brasil, depositados no herbário do Instituto de Botânica. Rickia 9: 37–53.

Gugliotta A, Bononi VLR. 1999. *Polyporaceae* do Parque Estadual da Ilha do Cardoso, São Paulo, Brasil. Bol. Inst. Bot. (São Paulo) 12: 1–112.

Gugliotta A, Capelari M. 1995. *Polyporaceae* from Ilha do Cardoso , SP, Brazil. Mycotaxon 56: 107–113.

Holmgren PK, Holmgren NH. 1998 Index Herbariorum: New York Botanical Garden's Virtual Herbarium. Available at: http://sweetgum.nybg.org/ih/Holmegren.

Jesus MA. 1993. Basidiomicetos lignocelulolíticos de floresta nativa e de *Pinus elliottii* Engelm. do Parque Estadual das Fontes do Ipiranga, São Paulo, SP. Hoehnea 20(1/2): 119–126.

Leal GR, Gugliotta AM. 2008. Criptógamos do Parque Estadual das Fontes do Ipiranga, São Paulo, SP. Fungos, 9: *Meripilaceae*. Hoehnea 35(1): 99–110.

Louza GSG, Gugliotta AM. 2007. *Polyporus* Fr.(*Polyporaceae*) no Parque Estadual das Fontes do Ipiranga, São Paulo, SP, Brasil. Hoehnea 34(3): 365–382.

Mittermeier RA, Myers N, Gil PR, Mittermeier CG. 1999. Hotspots: earth's biologically richest and endangered terrestrial ecoregions. Mexico, CEMEX. 430 p.

Secretaria de Estado de Meio Ambiente. 1996. Atlas das Unidades de Conservação Ambiental do Estado de São Paulo – parte I litoral. São Paulo, SMA. 30 p. + 7 mapas.

Secretaria de Estado de Meio Ambiente. 2000. Atlas das Unidade de Conservação Ambiental do Estado de São Paulo. São Paulo, SMA. 64 p. + 19 mapas.

Soares SCS, Gugliotta AM. 1998. Criptógamos do Parque Estadual das Fontes do Ipiranga, São Paulo, SP. Fungos, 7: *Aphyllophorales* (*Hymenochaetaceae*). Hoehnea 25(1): 11–31.

SOS Mata Atlântica, Instituto Nacional de Pesquisas Espaciais. 2009. Atlas dos remanescentes florestais da Mata Atlântica, Período 2000 a 2005. <http://www.sosma.org.br>.